



SUPRALYT®

Diætetisk foder til stabilisering af væske- og elektrolytbalance /
Dietetic feed for stabilization of water and electrolyte balance

Sammensætning/Composition

Glucose, natriumbicarbonat, maltodextrin, sodiumchlorid, kaliumklorid/
Glucose, sodium bicarbonate, maltodextrose, sodium chloride,
potassium chloride.

Tørstof/Dry matter	%	98
Råprotein/crude protein	%	0,1
Råaske/Ash	%	20,4
Calcium/Ca	g/kg	0,6
Natrium/Sodium	g/kg	70
Natriumbicarbonat/Sodium Bicarbonate	g/kg	150
Klorid/Cl	g/kg	60
Fodersalt/NaCl	g/kg	70
Kalium/K	g/kg	17
Druesukker/Sugar	g/kg	630
Maltodextrin/Maltodextrose	g/kg	150

Tilsætningsstoffer/Feed additives

Vitamin A, 3a672a	IU/kg	100.000
Vitamin D3, 3a671	IU/kg	10.000
Vitamin E, 3a700 all-rac-alpha-tocopheryl acetate	mg/kg	350
Vitamin B1, thiamine mononitrate, 3a821	mg/kg	60
Vitamin B2, riboflavin, 3a825i	mg/kg	82
Vitamin B6, 3a831 pyridoxine hydrochl.	mg/kg	62
Vitamin B12, 3a835 cyanocobalamin	mg/kg	0,2
D-Biotin, 3a880	mg/kg	4
Folinsyrefolc acid, 3a316	mg/kg	30
Niacin, 3a315 niacinamide	mg/kg	600
Pantothenic acid, calcium D-pantothen, 3a841	mg/kg	300
Choline, 3a890 choline chloride	mg/kg	520
Vitamin C, 3a300 L-ascorbic acid	mg/kg	4000

Tilskudsfoder

Complementary
feed

10 kg

Holdbarhed/Shelf life: 18 måneder/month.

Opbevaring/storage conditions: Rumtemperatur og beskyttet mod fugt/Room temperature and protected from moisture.

Brugsanvisning og dosering:

Supralyt anvendes til kalve, svin, lam, kid og føl, hvis der foreligger risiko for fordøjelsesforstyrrelser og i perioder med fordøjelsesforstyrrelser. Bland 175 g Supralyt i 5 liter varmt vand (40 - 50 gr. C).

Anvendes i 1 - 7 dage efter behov.

Supralyt må maximum anvendes som eneste foder i 2 dage.

Når Supralyt tildeles forebyggende, anvendes halv dosis.

Rengør og desinficer drikketrug dagligt.

1 dl svarer til ca 90 g.

Det anbefales at indhente råd fra en dyrlæge inden brug eller inden forlængelse af brugsperioden.

Fremstillet af: aFl21110001.

Dosage and use instruktion:

Calves, pigs, lambs and foal, in case of risk and during periods of diarrhea and recovery from digestive disturbance. Mix 35 g in 1 liter of warm water (40-50°C).

Can be used 1-7 days as needed. Supralyt must only be used as sole food for 2 days. If used preventive use half dosage.

1 dl corresponds to approx 90 g.

It is recommended to follow veterinarian instruction first.

Manufactured of: aFl21110001.



Partinummer / Lot nr:

Mindst holdbar til / Best before:

vifofarm
+45 8887 5300